

LETTERS TO THE EDITOR

## Pulmonary artery dilatation and obstructive sleep apnea

Joseph Simonson, MD<sup>1</sup>; Harley Greenberg, MD, FCCP<sup>1,2</sup>; Arunabh Talwar, MD, FCCP<sup>1,2</sup>

<sup>1</sup>Northwell Health, Division of Pulmonary Medicine, Critical Care and Sleep Medicine, New Hyde Park, New York; <sup>2</sup>Heart and Lung Research Unit, The Feinstein Institute for Medical Research, Manhasset, New York

We read with great interest the article by Nokes et al<sup>1</sup> reporting a high incidence of sleep-disordered breathing [SDB] in patients with scleroderma as evaluated by overnight pulse oximetry. It is interesting to note that the authors report in their study that in scleroderma patients, the only predictive variables for abnormal overnight oximetry [defined as [ODI >5/h or a mean oxygen saturation < 90%] were advancing age and pulmonary artery diameter >3.0 cm.

Pulmonary artery dilatation can occur from many causes, including abnormal pulmonary hemodynamics, hypoxemia, trauma, and vasculitis.<sup>2-4</sup> During last few years, there has been significant improvement in the understanding of the link between SDB, pulmonary hypertension,<sup>5</sup> and interstitial lung diseases.<sup>6</sup>

There is also a suggestion that obstructive sleep apnea is associated with enlargement of the main pulmonary artery.<sup>7,8</sup> The current study,<sup>1</sup> although retrospective, again highlights that in patients with SDB, repetitive nocturnal hypoxemia, and large intrathoracic negative pressure swings attributable to recurrent upper airway obstruction could affect pulmonary hemodynamics and may lead to pulmonary artery dilatation, although we agree that this concept needs to be studied further.

Patients with scleroderma report features of sleepiness and fatigue, usually attributed to the scleroderma disease process itself rather than to underlying SDB. As clinicians, we believe the presence of an enlarged pulmonary artery in scleroderma patients should not only rule out pulmonary hypertension but also remind ourselves about the possibility of underlying SDB.

### CITATION

Simonson J, Greenberg H, Talwar A. Pulmonary artery dilatation and obstructive sleep apnea. *J Clin Sleep Med*. 2020; 16(4):647.

### REFERENCES

1. Nokes BT, Raza HA, Cartin-Ceba R, et al. Individuals with scleroderma may have increased risk of sleep-disordered breathing. *J Clin Sleep Med*. 2019;15(11):1665–1669.
2. Raymond TE, Khabbaza JE, Yadav R, Tonelli AR. Significance of main pulmonary artery dilation on imaging studies. *Ann Am Thorac Soc*. 2014;11(10):1623–1632.
3. McCall RK, Ravenel JG, Nietert PJ, Granath A, Silver RM. Relationship of main pulmonary artery diameter to pulmonary arterial pressure in scleroderma patients with and without interstitial fibrosis. *J Comput Assist Tomogr*. 2014;38(2):163–168.
4. Bakker E, Ninaber M, Stolk J. Pulmonary artery dilation and increased lung density in systemic sclerosis. *Eur Respir J*. 2015;46:PA3743.
5. Minic M, Granton JT, Ryan CM. Sleep disordered breathing in group 1 pulmonary arterial hypertension. *J Clin Sleep Med*. 2014;10(3):277–283.
6. Kim JS, Podolanczuk AJ, Borker P, et al. Obstructive sleep apnea and subclinical Interstitial Lung Disease in the Multi-Ethnic Study of Atherosclerosis (MESA). *Ann Am Thorac Soc*. 2017;14(12):1786–1795.
7. Yakut T, Balcan B, Karakurt S. Determinants of enlarged main pulmonary artery diameter in patients with scleroderma and concomitant obstructive sleep apnoea. *Eur Respir J*. 2018;52:PA429.
8. Dobrowolski P, Florczak E, Klisiewicz A, et al. Pulmonary artery dilation indicates severe obstructive sleep apnea in patients with resistant hypertension: the Resist-POL Study. *Pol Arch Med Wewn*. 2016;126:222–229.

### SUBMISSION & CORRESPONDENCE INFORMATION

Submitted for publication December 9, 2019

Submitted in final revised form December 12, 2019

Accepted for publication December 12, 2019

Address correspondence to: Arunabh Talwar, MD, FCCP, Northwell Health, Pulmonary, Critical Care and Sleep Medicine, 410 Lakeville Rd. New Hyde Park, NY 11040, Tel: (516) 465-5400, Fax: (516) 465-5454, E-mail: arunabh@northwell.edu

### DISCLOSURE STATEMENT

On behalf of all the contributors Arunabh Talwar, MD is the guarantor of the data for this manuscript. All authors have seen and approved the manuscript. The authors report no conflicts of interest.