

GABA_A Receptors: Subtypes, Regional Distribution, and Function

David Nutt, D.M.

Psychopharmacology Unit, University of Bristol School of Medicine Sciences, Bristol, United Kingdom

Abstract: Modulatory agents of gamma-aminobutyric acid type A (GABA_A) receptors have been widely used for more than 40 years to treat anxiety, epilepsy, and sleep disorders; these drugs are generally safe, well tolerated, and effective. Recently, there has been a substantial growth in understanding of the mechanism of action of these drugs, which act at different sites on GABA_A receptors. A variety of GABA_A receptor subtypes with distinct functional roles have been characterized and an evolving

awareness of GABA_A receptor modulation holds promise for the future development of new, more sophisticated drug interventions that can elicit more selective effects by targeting specific subtypes of GABA_A receptors. Advances in genetic engineering have led to the development of transgenic mouse models that have further refined our understanding of the pharmacology and physiology for various GABA_A receptor subunits.

Gamma-aminobutyric acid (GABA) is the most widely distributed inhibitory neurotransmitter in the central nervous system (CNS).¹ As such, GABA limits the excitability of neuronal activity in all areas of the brain.² Excessive GABAergic signaling results in sedation, amnesia, and ataxia, whereas the mildest attenuation of GABAergic signaling results in arousal, anxiety, restlessness, insomnia, and exaggerated reactivity.

GABA is synthesized in presynaptic neurons and stored in synaptic vesicles; the rate-limiting step in the process is modulated by the activity of L-glutamic acid decarboxylase, an enzyme that promotes the synthesis of GABA.³ Upon neuronal activation, GABA is released from vesicles into the synapse, where it can act on postsynaptic receptors, or diffuse into the extracellular space and activate extrasynaptic receptors on postsynaptic neurons. GABA is then removed from the extracellular space by GABA transporters located on presynaptic neurons and glia that take up GABA into cells away from postsynaptic GABA receptors.^{4,5} Extracellular levels of GABA are regulated by the balance between GABA release from presynaptic vesicles and GABA uptake by GABA transporters.⁵

There are two main types of GABA receptors: fast-acting ionotropic GABA_A and GABA_C receptors, and slower-acting metabotropic GABA_B receptors.¹ GABA_A receptors are the predominant type of GABA receptor in the brain.⁶ It has been estimated that 20% to 50% of all central synapses contain ionotropic GABA_A receptors.²

Briefly, GABA_B receptors are metabotropic G-protein-linked receptors, coupled by intracellular signal transduction cascades to calcium and potassium channels. GABA_B receptors are the site of action of the muscle relaxant, baclofen, and are insensitive to drugs that modulate GABA_A receptors.¹ When GABA binds to the GABA_B receptor, there is an increase in potassium conductance

and a decrease in voltage-dependent calcium currents, resulting in hyperpolarization of the neuron and inhibition of neurotrans-




Figure 1a—The GABA_A receptor complex, visualized by electron microscopy, showing five protein subunits arranged around a central pore. Reproduced with permission from Nayeem N et al, *J Neurochem.* 1994;62:815-8.⁸

Figure 1b—Schematic representation of agonist binding sites and extracellular modulatory domains within the GABA_A receptor. Reproduced with permission from Whiting PJ et al. *Ann NY Acad Sci.* 1999;868:655-53.⁹

Disclosure Statement

Dr. Nutt has received research support from MSD, AZ, and Wyeth; has provided consulting services to MSD, Pfizer, Inc., Wyeth, and Organon; is a stock shareholder of GlaxoSmithKline; and is a member of the speakers' bureau for GlaxoSmithKline, Janssen, and Wyeth.




Figure 2a—Dendrogram depicting the known GABA_A receptor subunits and their sequence homologies grouped in clusters.

Figure 2b—The most prevalent GABA_A receptor in the central nervous system is the α₁β₂γ₂ receptor, depicted here with the appropriate stoichiometry of 2:2:1.

Figure 2c—A pie chart representing the approximate abundance of GABA_A receptor subtypes in the rat brain. Subscript x indicates the type of α, β or γ subunit is not known. This is a broad representation, and other receptor subunit combinations probably exist. Reproduced with permission: 2a⁴; 2b²; 2c¹¹.

mission.⁷

GABA_A receptors consist of five protein subunits arranged in a ring around a central pore; upon activation, the receptor allows chloride ions to flow into the cell, (Figure 1a)^{2,8} resulting in neuronal hyperpolarization.²

Benzodiazepines enhance the effects of GABA by lowering the concentration of GABA required to open the GABA_A channel.² Benzodiazepines bind to a modulatory site on the GABA_A receptor that is distinct from the GABA binding site (Figure 1b)⁹ and allosterically change the receptor complex to increase the affinity of the receptor to GABA, producing a larger post-synaptic current, prolonging inhibition.¹⁰ Although benzodiazepines do not act directly to open the chloride channel, they modulate the ability of GABA to do so, thus augmenting its inhibitory effects.

In addition to GABA and benzodiazepines, additional psychoactive compounds, such as barbiturates, loreclezole, picrotoxin, and neurosteroids can also bind to the GABA_A receptor and directly open the chloride channel (Figure 1b).⁹ Drugs such as barbiturates, chloral hydrate, clomethiazole, and ethanol enhance GABAergic transmission but at high concentrations can directly open the chloride channel. Because of this direct action on the chloride channel, these drugs have a propensity to be fatal in overdose.²

Heterogeneity of GABA_A Receptor Subtypes

There is a high level of heterogeneity of functional GABA_A receptors. GABA_A receptors are formed from co-assembly of at least 19 different subunits encoded by different genes (Figure 2a).^{4,11} These subunits are grouped into α, β, γ, δ, ε, and σ subunits, according to sequence homology; δ, ε, and σ subunits are less common and their functions have not yet been fully determined. GABA_A receptor subunits assemble into a limited number of pentameric receptors with the general stoichiometry of 2 α, 2 β and 1 γ subunits; δ and ε subunits may substitute for the γ subunit, and the σ subunit may substitute for the β subunit (Figure 2b).² The majority of GABA_A receptors expressed in the CNS are α₁β₂γ₂; with α₃β₃γ₂ and α₂β₃γ₂ receptors also highly prevalent (Figure 2c).¹¹ GABA binds at the junction between subunits α and β, whereas benzodiazepines bind at the interface between subunits α and γ (Figure 1b).⁹

Immunocytochemistry data suggest that there is high expression of GABA_A receptor subunits α₁, β₁, β₂, β₃, and γ₂ throughout the brain, though there are differences in their distribution.¹² The most widely expressed subunits are the α₁ and γ₂ subunits and these subunits are therefore likely constituents of a high proportion of GABA_A receptors. The α₂, α₃, α₄, α₅, α₆ subunits have more confined distributions, which do not often overlap. For example, α₅ subunits are highly localized in the olfactory bulb, hypothalamus, and the hippocampus.¹² The α₄ subunits are concentrated in the thalamus, hippocampus, olfactory tubercle, and basal ganglia. The distribution of the δ subunit parallels that of the α₄ subunits, in the thalamus, striatum, outer layers of the cortex, and the hippocampus.¹² Similar findings in subunit distribution have been demonstrated using in situ hybridization to measure expression levels of mRNA encoding the different GABA_A receptor subunits.¹³

Further specificity of GABA_A receptor expression exists among different neuronal cell types. For example, a neuron-specific pattern of subunit distribution is apparent when the α-subunits are co-visualized in double immunofluorescence staining with mark-

ers for GABAergic, catecholaminergic, or serotonergic neurons. Neurons within the pallidum express only α_1 subunits, whereas the brainstem reticular formation expresses both α_1 and α_3 subunits. An additional population of α_3 -containing, but not α_1 -containing, neurons is seen in monoaminergic neurons.¹⁴ Furthermore, most serotonergic neurons in the raphe nuclei express high levels of only the α_3 subunit, but not α_1 subunits; in contrast, GABAergic neurons express both α_1 and α_3 subunits.¹⁵ Thus, great heterogeneity of both subunit type and expression patterns of GABA_A receptors occurs throughout the CNS.

GABA_A receptor subunits are also differentially expressed within specific subcellular regions within individual neurons, and such selective trafficking and localization of GABA_A receptor subunits contributes to the functional properties of GABA_A inhibition.^{16,17} For example, within the granule cells of the cerebellum, α_1 , α_6 , $\beta_{2/3}$, and γ_2 subunits are localized in high concentrations at Golgi cell synapses and in lower concentrations within the extrasynaptic membrane. In contrast, δ subunits are absent from synaptic junctions but are abundant in the extrasynaptic dendritic and somatic membranes. In addition, α_6 , γ_2 , and $\beta_{2/3}$ subunits, but not the δ subunits, are also concentrated in some glutamatergic mossy fiber synapses.¹⁸ This pattern of subcellular localization has important implications regarding receptor function. The δ -subunit-containing receptors are expressed extrasynaptically, and are thought to play a role in the GABA-mediated overall excitability of postsynaptic neurons through tonic inhibition.¹¹ Such GABA-mediated tonic inhibition has been observed in granule cells of the cerebellum, dentate gyrus, and in hippocampal interneurons and is thought to occur as a result of extrasynaptic GABA_A receptor activation by spillover of GABA from the synapse.^{19,20}

Diverse Functions of GABA_A Receptor Subunits

Although GABA_A receptor subtypes have distinct anatomical and subcellular localizations, it is now known that only a subset of GABA_A receptors mediate the effects of benzodiazepines: $\alpha_1\beta\gamma_2$ -, $\alpha_2\beta\gamma_2$ -, $\alpha_3\beta\gamma_2$ -, and $\alpha_5\beta\gamma_2$ -containing receptors bind benzodiazepines with high affinity, but $\alpha_4\beta\gamma_2$ - and $\alpha_6\beta\gamma_2$ -containing receptors have a lower affinity for benzodiazepines.¹¹

The function of specific GABA_A receptor subunits has been unraveled further by pharmacological and behavioral studies of transgenic mice in which different GABA_A receptor subunits are either deleted due to gene ablation (knockout [KO] mice) or mutated in a manner that creates a nonfunctional receptor site (knock-in mice) (Table 1). Several studies have examined the contribution of different GABA_A receptor subunits to the effects of benzodiazepines. Transgenic mouse lines with mutations postulated to alter individual amino acid residues in the benzodiazepine binding site have revealed that a single amino acid is responsible for high-affinity benzodiazepine binding. This is the histidine residue at position 101 of the primary amino acid sequence of the α_1 subunit and at equivalent positions of the sequences of α_2 , α_3 , and α_5 . Low affinity binding is conferred by an arginine in the equivalent position in the protein structure of the α_4 and α_6 subunits. Knock-in mice, containing a mutation to this requisite residue, are insensitive to benzodiazepines but remain responsive to GABA.²¹

Studies of knock-in mice that express a benzodiazepine-insensitive α_1 subunit reveal an important role of α_1 subunits in benzodiazepine-induced sedation. These knock-in mice fail to show the sedative, amnesic effects of diazepam, and the anticonvulsant

Table 1—The Roles of Select GABA_A Receptor Subunits²¹

	α_1	α_2	α_3	α_5	γ_2	β_2	β_3	δ
Effects of benzodiazepines:								
Sedation	+	-	-	-		+		
Anxiolysis	-	+	-/+	-				
Amnesia	+			+				
Myorelaxation	-		+					
Motor impairment	-	-	-					
Anticonvulsant	+	-	-	-				
Ethanol reinforcement	-			+				
Effects of anesthetics								
Anxiety	+					+	+	+
Learning/memory				+				+

Adapted from Rudolph U, 1999.²¹

effects of diazepam are partly reduced. These mice do retain other effects of diazepam, including the anxiolytic-like, myorelaxant, motor-impairing, and ethanol-potentiating actions.²¹ These results suggest that sedative, amnesic, and anticonvulsant effects of diazepam are mediated in part by wild-type GABA_A receptor subunits, namely the α_2 , α_3 , and α_5 subunits.²¹ The results of this study are supported by other studies that demonstrate that mice with this specific mutation in the α_1 subunit fail to exhibit the sedative effect of benzodiazepines, but exhibit a normal anxiolytic response, equivalent to that of wild-type mice.²² An earlier study had created mice lacking the α_1 subunit, and these KO mice showed a marked decrease in sleep induced by nonselective benzodiazepine, flurazepam, and the α_1 -selective hypnotic drug zolpidem.²³ GABA_A receptor β_2 subunits may also play a role in sleep induced by benzodiazepines as the same paper demonstrates that β_2 -subunit KO mice show a marked decrease in sleep induced by the nonselective benzodiazepine, flurazepam, and the α_1 -selective drug zolpidem.²³ Together, these results suggest that a drug that acts as an agonist at α_2 , α_3 , or α_5 subunit-containing GABA_A receptors may provide anxiolysis without inducing sedation.

The anxiolytic effect of benzodiazepines appears to be attributable to the α_2 and/or the α_3 subunit of the GABA_A receptor. Transgenic mice that express a mutated form of the α_2 subunit, in which the histidine at residue 101 is replaced by arginine, fail to exhibit an anxiolytic response to treatment with diazepam.²⁴ Diazepam has an anxiolytic effect in wild-type and α_3 -subunit KO mice, but not in α_2 -subunit KO mice, whereas the sedative, motor-impairing, and anticonvulsant actions of diazepam are not impaired in the α_2 - or α_3 -subunit KO mice.²⁴

There is more evidence to support the role for α_2 and α_3 subunits, but not α_1 subunits, in anxiolysis. A novel compound, L838,417, is an antagonist at α_1 -containing GABA_A receptors, but has agonist activity at α_2 -, α_3 -, and α_5 -containing receptors, and is a nonsedating anxiolytic. Whether administered to either wild-type mice or α_1 -subunit KO mice, L838,417 has an equivalent anxiolytic effect.²²

Another GABA_A receptor subunit that may play a role in anxiolytic pharmacology is the γ_2 subunit. Transgenic mice that express lower levels of the γ_2 subunit through KO of one of the two alleles display enhanced anxiety-like behavior on the elevated plus maze and forced novel exploratory tests. These mice have unaltered hypnotic response to benzodiazepines.²⁵

Other studies in transgenic mouse lines have also elucidated the roles of different GABA_A receptor subunits in anesthesia.




Figure 3—Pseudocolor PET image represents a global decrease in benzodiazepine binding sites in patients with panic disorder (right) versus healthy control patients (left). Reproduced with permission from Tiihonen J.³⁷

Male, but not female, α_1 - and β_2 -subunit KO mice exhibit reduced ethanol-induced sleep, whereas β_2 - (but not α_1 -) KO mice of both sexes show diminished sleep in response to the β -selective anesthetic etomidate. These mice showed normal responses to pentobarbital, suggesting that pentobarbital's effects at other ion channels may be crucial to its remaining effects on the CNS.²³

Another study found that α_1 -subunit KO mice exhibit normal ethanol-induced anxiolysis, ataxia, anticonvulsant, and hypnotic effects. The anticonvulsant effects of neurosteroids were also unaltered in these mice. However the α_1 -subunit KO mice did exhibit reduced hypnotic response with pentobarbital, etomidate, and midazolam as well as showing an enhanced responses to ketamine.²⁶ In another study, a mouse line with a mutation of a single residue of the protein sequence of the β_3 subunit of the GABA_A receptor failed to exhibit anesthetic response to treatment with etomidate and propofol, and exhibited slightly decreased responses to the volatile anesthetics enflurane and halothane.²⁷ In addition, the same strain of mice require higher isoflurane levels for anesthesia.²⁸ Mutant mice, which do not express α_6 subunits, show normal response to various general anesthetics.²⁹ However, mice lacking the δ subunit show attenuated sensitivity to this action of neurosteroids, but normal response to etomidate, propofol, and halothane.³⁰ Thus, whereas anesthetics in general appear to be active at the β_3 subunit, volatile anesthetics may have additional sites of action that have yet to be clearly characterized.

Other interesting findings have been derived from studies of GABA_A receptor subunit KO mouse lines. Mice devoid of the GABA_A receptor β_3 subunit exhibit spontaneous epilepsy and hyperactivity, supporting a link between GABA neurotransmission and epilepsy.³¹ Studies of two transgenic mouse lines reveal an important role of α_5 subunits in learning and memory. These include a KO mouse line, which does not express the α_5 subunit,³² and a mouse line in which the histidine-arginine mutation has been introduced.³³ This point mutation leads to a significant decrease in the expression of the α_5 subunit in the hippocampus, a region of the brain that is known to have an important role in learning and memory.³⁴ Both lines of these transgenic mice show improved performance in animal models of learning and memory.^{32,33} These results suggest that a selective inhibitor of α_5 -containing GABA_A receptors could have use as a cognitive enhancer to treat mild

cognitive impairment in the elderly or in patients with Alzheimer's disease. Another KO mouse line, devoid of δ subunits of the GABA_A receptor, also demonstrates enhanced performance in a model of learning and memory; compared with controls, female δ -subunit KO mice exhibited enhanced conditional acquisition of tone and context fear.³⁵

In Vivo Imaging of Human GABA_A Receptors

A novel technique that offers a promising method for characterizing the roles of GABA_A receptor subtypes in normal physiology and in pathologic disease is the in vivo imaging of GABA_A receptor subunits in the human brain with PET (positron emission tomography) or SPECT (single photon emission computed tomography). PET imaging has been used to measure GABA_A receptor quantity and localization within human subjects using a radiolabeled benzodiazepine-site antagonist, flumazenil. A PET study conducted in patients with panic disorder found a global reduction in GABA_A receptor density throughout the brain when compared with control subjects (Figure 3). The loci with the largest regional decrease in binding, the right orbitofrontal cortex and right insula, are areas thought to be essential in the central mediation of anxiety.³⁶ Other studies have shown more circumscribed reductions in GABA_A receptor binding in two other anxiety disorders—generalized anxiety disorder (GAD)³⁷ and post-traumatic stress disorder (PTSD)³⁸—suggesting reduced central inhibition may be a common feature of anxiety disorders.

With the development of GABA_A receptor subunit-selective tracers, the roles of the subunits may be examined in vivo in healthy humans and in patients with a variety of disorders. The α_5 subunit-selective agent Ro15-4513 can be used to selectively visualize α_5 subunits in humans by PET, and can lead to further exploration of the functional role of this GABA_A receptor subtype in humans.³⁹

CONCLUSIONS

How to Exploit the Heterogeneity of GABA_A Receptors for Drug Development

The emerging understanding of the roles of GABA_A receptors can be utilized in various ways for drug development. Selective drugs can be developed to target specific GABA_A receptor subunits implicated in specific functions. For example, zolpidem is a GABA_A receptor α_1 subunit-selective compound, indicated for the short-term treatment of insomnia. Furthermore, gaboxadol (THIP, 4,5,6,7-tetrahydroisoxazolo(5,4-c)pyridin-3-ol) is an extrasynaptic GABA_A-selective agonist that primarily targets benzodiazepine-insensitive $\alpha_4\beta_3\delta$ receptors and is in the late stages of clinical development for the treatment of insomnia.⁴⁰ Other potentially useful drugs may be α_2 subunit- or α_3 subunit-selective agonists as nonsedating anxiolytics, and α_5 subunit inverse agonists as memory enhancers.^{41,42} Future research will further elucidate the roles of GABA_A receptor subunits and provide applications for this newfound knowledge.

REFERENCES

1. Bormann J. The 'ABC' of GABA receptors. *Trends Pharmacol Sci.* 2000;21:16-9.
2. Nutt DJ, Malizia AL. New insights into the role of the GABA(A)-benzodiazepine receptor in psychiatric disorder. *Br J Psychiatry.*

- 2001;179:390-6.
3. Jin H, Wu H, Osterhaus G et al. Demonstration of functional coupling between gamma-aminobutyric acid (GABA) synthesis and vesicular GABA transport into synaptic vesicles. *Proc Natl Acad Sci USA*. 2003;100:4293-8.
 4. Cherubini E, Conti F. Generating diversity at GABAergic synapses. *Trends Neurosci*. 2001;24:155-62.
 5. Deken SL, Wang D, Quick MW. Plasma membrane GABA transporters reside on distinct vesicles and undergo rapid regulated recycling. *J Neurosci*. 2003;23:1563-8.
 6. Bowerly NG, Hudson AL, Price GW. GABAA and GABAB receptor site distribution in the rat central nervous system. *Neuroscience*. 1987;20:365-83.
 7. Ong J, Kerr DI. Recent advances in GABAB receptors: from pharmacology to molecular biology. *Acta Pharmacol Sin*. 2000;21:111-23.
 8. Nayeem N, Green TP, Martin IL, Barnard EA. Quaternary structure of the native GABAA receptor determined by electron microscopic image analysis. *J Neurochem*. 1994;62:815-8.
 9. Whiting PJ, Bonnert TP, McKernan RM et al. Molecular and functional diversity of the expanding GABA-A receptor gene family. *Ann N Y Acad Sci*. 1999;868:645-53.
 10. Smith GB, Olsen RW. Functional domains of GABAA receptors. *Trends Pharmacol Sci*. 1995;16:162-8.
 11. Whiting PJ. GABA-A receptor subtypes in the brain: a paradigm for CNS drug discovery? *Drug Discov Today*. 2003;8:445-50.
 12. Pirker S, Schwarzer C, Wieselthaler A, Sieghart W, Sperk G. GABA(A) receptors: immunocytochemical distribution of 13 subunits in the adult rat brain. *Neuroscience*. 2000;101:815-50.
 13. Wisden W, Laurie DJ, Monyer H, Seeburg PH. The distribution of 13 GABAA receptor subunit mRNAs in the rat brain. I. Telencephalon, diencephalon, mesencephalon. *J Neurosci*. 1992;12:1040-62.
 14. Fritschy JM, Benke D, Mertens S, Oertel WH, Bachi T, Mohler H. Five subtypes of type A gamma-aminobutyric acid receptors identified in neurons by double and triple immunofluorescence staining with subunit-specific antibodies. *Proc Natl Acad Sci USA*. 1992;89:6726-30.
 15. Gao B, Fritschy JM, Benke D, Mohler H. Neuron-specific expression of GABAA-receptor subtypes: differential association of the alpha 1- and alpha 3-subunits with serotonergic and GABAergic neurons. *Neuroscience*. 1993;54:881-92.
 16. Loup F, Weinmann O, Yonekawa Y, Aguzzi A, Wieser HG, Fritschy JM. A highly sensitive immunofluorescence procedure for analyzing the subcellular distribution of GABAA receptor subunits in the human brain. *J Histochem Cytochem*. 1998;46:1129-39.
 17. Nusser Z, Sieghart W, Benke D, Fritschy JM, Somogyi P. Differential synaptic localization of two major gamma-aminobutyric acid type A receptor alpha subunits on hippocampal pyramidal cells. *Proc Natl Acad Sci USA*. 1996;93:11939-44.
 18. Nusser Z, Sieghart W, Somogyi P. Segregation of different GABAA receptors to synaptic and extrasynaptic membranes of cerebellar granule cells. *J Neurosci*. 1998;18:1693-703.
 19. Brickley SG, Cull-Candy SG, Farrant M. Development of a tonic form of synaptic inhibition in rat cerebellar granule cells resulting from persistent activation of GABAA receptors. *J Physiol*. 1996;497 (Pt 3):753-9.
 20. Kullmann DM, Ruiz A, Rusakov DM, Scott R, Semyanov A, Walker MC. Presynaptic, extrasynaptic and axonal GABAA receptors in the CNS: where and why? *Prog Biophys Mol Biol*. 2005;87:33-46.
 21. Rudolph U, Crestani F, Benke D et al. Benzodiazepine actions mediated by specific gamma-aminobutyric acid(A) receptor subtypes. *Nature*. 1999;401:796-800.
 22. McKernan RM, Rosahl TW, Reynolds DS et al. Sedative but not anxiolytic properties of benzodiazepines are mediated by the GABA(A) receptor alpha1 subtype. *Nat Neurosci*. 2000;3:587-92.
 23. Blednov YA, Jung S, Alva H et al. Deletion of the alpha1 or beta2 subunit of GABAA receptors reduces actions of alcohol and other drugs. *J Pharmacol Exp Ther*. 2003;304:30-6.
 24. Low K, Crestani F, Keist R et al. Molecular and neuronal substrate for the selective attenuation of anxiety. *Science*. 2000;290:131-4.
 25. Chandra D, Korpi ER, Miralles CP, De Blas AL, Homanics GE. GABAA receptor gamma2 subunit knockdown mice have enhanced anxiety-like behavior but unaltered hypnotic response to benzodiazepines. *BMC Neurosci*. 2005;6:30.
 26. Kralic JE, Wheeler M, Renzi K et al. Deletion of GABAA receptor alpha 1 subunit-containing receptors alters responses to ethanol and other anesthetics. *J Pharmacol Exp Ther*. 2003;305:600-7.
 27. Jurd R, Arras M, Lambert S et al. General anesthetic actions in vivo strongly attenuated by a point mutation in the GABA(A) receptor beta3 subunit. *FASEB J*. 2003;17:250-2.
 28. Liao M, Sonner JM, Jurd R et al. Beta3-containing gamma-aminobutyric acidA receptors are not major targets for the amnesic and immobilizing actions of isoflurane. *Anesth Analg*. 2005;101:412-8, table.
 29. Homanics GE, Ferguson C, Quinlan JJ et al. Gene knockout of the alpha6 subunit of the gamma-aminobutyric acid type A receptor: lack of effect on responses to ethanol, pentobarbital, and general anesthetics. *Mol Pharmacol*. 1997;51:588-96.
 30. Mihalek RM, Banerjee PK, Korpi ER et al. Attenuated sensitivity to neuroactive steroids in gamma-aminobutyrate type A receptor delta subunit knockout mice. *Proc Natl Acad Sci USA*. 1999;96:12905-10.
 31. Homanics GE, Delorey TM, Firestone LL et al. Mice devoid of gamma-aminobutyrate type A receptor beta3 subunit have epilepsy, cleft palate, and hypersensitive behavior. *Proc Natl Acad Sci USA*. 1997;94:4143-8.
 32. Collinson N, Kuenzi FM, Jarolimek W et al. Enhanced learning and memory and altered GABAergic synaptic transmission in mice lacking the alpha 5 subunit of the GABAA receptor. *J Neurosci*. 2002;22:5572-80.
 33. Crestani F, Keist R, Fritschy JM et al. Trace fear conditioning involves hippocampal alpha5 GABA(A) receptors. *Proc Natl Acad Sci USA*. 2002;99:8980-5.
 34. Burgess N, Maguire EA, O'Keefe J. The human hippocampus and spatial and episodic memory. *Neuron*. 2002;35:625-41.
 35. Wiltgen BJ, Sanders MJ, Ferguson C, Homanics GE, Fanselow MS. Trace fear conditioning is enhanced in mice lacking the delta subunit of the GABAA receptor. *Learn Mem*. 2005;12:327-33.
 36. Malizia AL, Cunningham VJ, Bell CJ, Liddle PF, Jones T, Nutt DJ. Decreased brain GABA(A)-benzodiazepine receptor binding in panic disorder: preliminary results from a quantitative PET study. *Arch Gen Psychiatry*. 1998;55:715-20.
 37. Tiihonen J, Kuikka J, Rasanen P et al. Cerebral benzodiazepine receptor binding and distribution in generalized anxiety disorder: a fractal analysis. *Mol Psychiatry*. 1997;2:463-71.
 38. Bremner JD, Innis RB, Southwick SM, Staib L, Zoghbi S, Charney DS. Decreased benzodiazepine receptor binding in prefrontal cortex in combat-related posttraumatic stress disorder. *Am J Psychiatry*. 2000;157:1120-6.
 39. Lingford-Hughes A, Hume SP, Feeney A et al. Imaging the GABA-benzodiazepine receptor subtype containing the alpha5-subunit in vivo with [11C]Ro15 4513 positron emission tomography. *J Cereb Blood Flow Metab*. 2002;22:878-89.
 40. Krosgaard-Larsen P, Frolund B, Liljefors T, Ebert B. GABA(A) agonists and partial agonists: THIP (Gaboxadol) as a non-opioid analgesic and a novel type of hypnotic. *Biochem Pharmacol*. 2004;68:1573-80.
 41. Chambers MS, Atack JR, Broughton HB et al. Identification of a novel, selective GABA(A) alpha5 receptor inverse agonist which enhances cognition. *J Med Chem*. 2003;46:2227-40.
 42. Chambers MS, Atack JR, Carling RW et al. An orally bioavailable, functionally selective inverse agonist at the benzodiazepine site of GABAA alpha5 receptors with cognition enhancing properties. *J Med Chem*. 2004;47:5829-32.