

Mood was assessed using Profile of Mood States (POMS); its 7 subscales (tension, anger, fatigue, depression, esteem-affect, vigor, confusion) were summed (with a constant of 100) to create a Total Mood Disturbance (TMD) score. Multiple linear regression models examined associations between sleep health and mood adjusting for age, gender, and whether pre- or post-COVID-19.

Results: The mean sleep health score was 4.7 ± 1.1 ; the mean TMD score was 96.6 ± 18.5 . Better sleep health was associated with lesser TMD ($\beta = -0.32$, $p < 0.001$) and better mood on each of the POMS subscales ($\beta \geq 0.18$, $p < 0.05$), aside from esteem-related affect ($p = 0.31$). Of the individual sleep dimensions, only satisfaction, alertness, and efficiency were associated with TMD ($\beta \geq 0.18$, $p < 0.05$). Satisfaction was the only individual sleep dimension that was consistently associated with better mood on each subscale ($\beta \geq 0.17$). Alertness, efficiency, and duration were inconsistently associated with individual mood subscales. Regularity and timing were not associated with any mood subscales ($p \geq 0.267$ and $p \geq 0.073$, respectively).

Conclusion: Better sleep health was associated with less TMD. Satisfaction was the sleep dimension that consistently associated with each subscale of mood. The cross-sectional, observational design limits casual inference between sleep health and mood disturbance due to a lack of temporality and the potential for residual confounding.

Support (If Any): This study was funded by National Institutes of Health (NIH) grants R01HL134809 and R01HL147610.

0254

ASSOCIATION OF SLOW WAVE ACTIVITY AND ODDS RATIO PRODUCT WITH INTERNALIZING AND EXTERNALIZING PROBLEMS IN CHILDREN AND ADOLESCENTS

Julio Fernandez-Mendoza¹, Anna Ricci¹, Fan He¹, Susan Calhoun¹, Jidong Fang¹, Magdy Younes², Alexandros Vgontzas¹, Duanping Liao¹, Edward Bixler¹

Penn State College of Medicine¹ University of Manitoba²

Introduction: The association of metrics of sleep microstructure with internalizing and externalizing problems in youth has remained elusive. While one study found increased frontal slow wave activity (SWA) in depressed adolescents, there is lack of evidence for a relationship between dimensional measures of behavior and metrics of sleep depth/intensity. We examined the association between two measures of sleep depth/intensity, slow wave activity (SWA) and odds ratio product (ORP), with internalizing and externalizing problems in children and adolescents.

Methods: We calculated SWA and ORP during non-rapid eye movement (NREM) sleep at central, frontal and fronto-occipital derivations in 639 children (5-12y, median 9y) and 418 adolescents (12-23y, median 16y) from the Penn State Child Cohort via in-lab polysomnography. ORP provides a standardized measure of NREM sleep depth, while ORP-9 (average ORP in the 9-seconds following NREM arousals) provides a metric of arousability. SWA (0.4-4Hz) absolute power (μV^2) was determined during NREM sleep. Internalizing and externalizing problems were assessed on Achenbach's Behavior Checklist by parent (subjects $\leq 17y$) or self-report (subjects $\geq 18y$). For each scale, T-scores with a mean of 50 and standard deviation of 10 were obtained following standardized scoring. Multivariable-adjusted linear regression models examined the association between SWA/ORP and clinical outcomes.

Results: At ages 5-12, fronto-occipital SWA was negatively associated with externalizing behaviors ($p = 0.05$), while fronto-occipital and frontal ORP, and frontal ORP-9 were positively associated with

internalizing symptoms (all $p < 0.01$). At ages 12-23, central SWA was negatively associated with internalizing symptoms ($p = 0.05$), while central ($p = 0.05$) and frontal ($p = 0.03$) ORP and central ORP-9 ($p = 0.03$) were positively associated with externalizing behaviors.

Conclusion: Reductions in SWA in childhood or adolescence are associated with developmentally appropriate behavioral problems, as depression/anxiety are more prevalent in adolescence. In contrast to SWA, increased ORP (lighter sleep) and ORP-9 (greater arousability) are associated with more anxiety/depression in childhood, yet more externalizing behaviors in adolescence. These distinct associations, such as SWA with externalizing behaviors and ORP with internalizing symptoms during childhood, may reflect how SWA captures local/synaptic control, while ORP global/state control, of sleep depth, making both sleep EEG biomarkers important from a developmental standpoint.

Support (If Any): National Institutes of Health (R01MH118308, UL1TR000127)

0255

SLEEP HYGIENE INDEX: ASSOCIATIONS WITH SLEEP AND MENTAL HEALTH IN COLLEGE STUDENTS

Matelyn Gibson¹, Noah Anderson¹, Alexis Horton¹, Kayla Mullins², Alexandria Reynolds¹

The University of Virginia's college at Wise¹ University of Tennessee²

Introduction: Typically, college students practice unhealthy sleep hygiene behaviors, obtain too little sleep, and experience poor sleep quality. Sleep hygiene includes the routines or practices that prepare a person for the best possible night of sleep. Good sleep hygiene habits, like creating a sleep-friendly environment and making time for sleep, promote healthy duration and quality of sleep. Stress is also an important factor to consider during the college experience. Sleep and mental health are tightly connected, and stress can negatively impact the sleep and mental health of individuals. The focus of the current study was to examine habitual sleep habits in college students, in association with sleep quality and psychological health.

Methods: Participants included 51 undergraduate students (18 men, average age $M = 20.25$ years, $SD = 1.78$) who wore wrist actigraphs to measure their typical sleep habits for one week. After one week, participants completed questionnaires about sleep quality (Pittsburgh Sleep Quality Index, PSQI) and sleep hygiene practices (Sleep Hygiene Index, SHI). Higher scores on PSQI represent poorer quality; higher scores on SHI represent unhealthy sleep hygiene behaviors. Mental health symptoms were measured by the Depression, Anxiety, and Stress Scale (DASS-21).

Results: Overall sleep duration was 6.59 hours and sleep efficiency was 82.55% as measured by actigraphy. PSQI scores ($M = 6.86$) demonstrated poor sleep quality and SHI scores ($M = 24.80$) indicated overall poor sleep hygiene practices. SHI scores predicted higher PSQI scores ($F(1, 50) = 18.05$, $p < .001$), but did not predict sleep duration or efficiency. Depression, anxiety, and stress scores on the DASS predicted poorer sleep hygiene ($F(1, 50) = 18.05$, $p < .001$; $F(1, 50) = 5.82$, $p = .020$; $F(1, 50) = 13.42$, $p < .001$; respectively).

Conclusion: As expected, college students' sleep was short in duration, poor in efficiency, and poor in quality. Additionally, poor sleep hygiene practices predicted poorer sleep quality. Interestingly, scores that indicated worse depression, anxiety, and stress predicted poorer sleep hygiene practices, suggesting that mental health may contribute to healthy sleep practices. More research is needed to understand the complex relationship between mental health, sleep, and healthy sleep practices

Support (If Any): None.