



ORIGINAL ARTICLE

Sleep disturbances among women in a Subarctic region: a nationwide study

Anna Bára Unnarsdóttir^{1,*}, Arna Hauksdóttir¹, Thor Aspelund^{1,6},
Vigdís Gunnarsdóttir¹, Gunnar Tómasson^{1,2}, Jóhanna Jakobsdóttir¹,
Unnur Anna Valdimarsdóttir^{1,3,4} and Edda Björk Thordardóttir¹

¹Centre of Public Health Sciences, Faculty of Medicine, School of Health Sciences, University of Iceland, Reykjavik, Iceland, ²Department of Rheumatology, University Hospital, Reykjavik, Iceland, ³Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden and ⁴Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, MA, USA

*Corresponding author. Anna Bára Unnarsdóttir, Centre of Public Health Sciences, Faculty of Medicine, University of Iceland, Sturlugata 8, 101 Reykjavik, Iceland. Email: abu@hi.is.

Abstract

Study Objectives: To date, few studies have assessed sleep problems among women residing in Subarctic regions. Therefore, the aim of this large-scale population-based study was to assess the prevalence of severe sleep problems and associated factors among Icelandic women, living at 63–66°N.

Methods: Participants were 29 681 women (18–69 years old) who took part in the Icelandic Stress-And-Gene-Analysis study in 2018–2019. Background information, health-related behavior, and mental health symptoms were assessed with an online questionnaire. The Pittsburgh Sleep Quality Index (PSQI) was used to assess severe sleep problems during the past month. Adjusting for age, marital status, number of children, education, personal income, work schedule, region, and response period, we used modified Poisson log-linear models to obtain prevalence ratios (PRs) with 95% confidence intervals (CIs).

Results: Overall, 24.2% of women reported severe sleep problems (PSQI >10). Women responding in the winter presented with an overall higher prevalence of severe sleep problems, compared to those responding in the summer (PR 1.21; 95% CI, 1.15 to 1.28). Severe sleep problems were more prevalent among young and late-midlife women, those who were single, had children, socio-economic challenges, worked shifts, and flexible hours. Furthermore, obesity, suboptimal health behaviors, excessive screen time, and mental health problems were associated with severe sleep problems.

Conclusion: Severe sleep problems are more common among women in Subarctic regions than elsewhere, particularly during winter. These findings motivate the development of preventive strategies and interventions for women in the Subarctic who suffer from sleep problems.

Statement of Significance

Limited studies have assessed sleep problems among women living in Subarctic regions, areas with extreme variation in seasonal light duration, with long dark periods during winter and constant daylight during summer. Our population-based study of 29 681 women in Iceland indicates that severe sleep problems are more prevalent among women responding during the winter months when daylight is limited. We also found socioeconomic challenges, shift work, and suboptimal health behaviors to be associated with higher prevalence of sleep problems among women in the Subarctic region of Iceland with high-social welfare. These results are valuable for identifying women with higher likelihood of sleep disruptions and therefore would benefit from targeted prevention, in the Subarctic.

Key words: sleep; PSQI; Subarctic; Iceland; women; population; seasons

Submitted: 29 December, 2021; Revised: 31 March, 2022

© Sleep Research Society 2022. Published by Oxford University Press on behalf of the Sleep Research Society. <https://creativecommons.org/licenses/by/4.0/>, which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

Introduction

Sleep is essential for human's overall well-being and general health [1], with most healthy adults needing at least 7 h of sleep each night to function properly and to avoid sleep deprivation [2]. Studies on sleep quality include both quantitative aspects, such as sleep length, sleep latency, and number of awakenings during the night, as well as qualitative aspects, such as feeling of being energetic and restored upon awakening [3]. Poor sleep quality has been associated with adverse long-term health consequences [4, 5], as well as higher risk of all-cause mortality [6].

Previous population-based studies indicate that 27%–52% of women experience sleep problems (Pittsburgh Sleep Quality Index [PSQI] > 5) [7–12] and 11%–28% severe sleep problems (PSQI > 10) [8, 13, 14]. Evidence suggests that insomnia, defined as having difficulties falling asleep or staying asleep [15], is more common among women in Subarctic regions (the Northern hemisphere), with 8%–16% of women suffering from insomnia in these regions [16–18], compared to 7%–12% women globally [19]. Winters in the Subarctic regions consist of dark periods while summers are marked by constant daylight [20]. It has been suggested that these light exposure patterns cause abnormalities of circadian rhythms and disturbed sleep [21–23]. Indeed, individuals who live near the Subarctic experience greater seasonal changes in insomnia and fatigue, compared to those who live near the equator [21]. However, few population-based studies have assessed sleep problems among Subarctic women living with extreme variation in seasonal light duration.

Previous results indicate that higher age [17], being in a relationship [24, 25], and having children [26, 27] are associated with poor sleep quality. Furthermore, previous studies have found low socio-economic status (SES) [7, 8, 28, 29] and shift work [30, 31] to be associated with sleep problems. In addition, high body mass index (BMI) [8, 32, 33] and suboptimal health behaviors, such as cigarette smoking [8, 9, 34, 35], excessive alcohol consumption [36, 37], and prolonged screen time [38–40] have been associated with sleep problems.

Sleep problems are also highly comorbid with many major mental disorders. It has been suggested that sleep problems are a contributory factor in the occurrence of disorders such as anxiety and depression [41]. However, few studies have assessed the degree to which these situational and health-related factors impact sleep among those residing in the Subarctic region.

There are some indications that sleep deprivation may be a problem in Iceland, with a recent report finding 24% of Icelandic women to sleep on average <6 h per night [42]. Dispensing of hypnotics and sedatives, used in the treatment for insomnia, are also considerably higher in Iceland than in other Nordic countries [43].

To the best of our knowledge, no population-based study has to date assessed the prevalence of sleep problems among women residing in these Subarctic regions and associated factors. Epidemiological studies with representative general population-based samples using validated measurements are needed. Therefore, the aim of the current study was to leverage a population-based cohort of Icelandic women to investigate the prevalence of sleep problems with regard to season and latitude of residency. Furthermore, to assess the association of severe sleep problems and demographic- and socioeconomic characteristics, as well as health behaviors.

Methods

Study population

Participants were women who took part in the baseline assessment of the population-based SAGA (Stress-And-Gene-Analysis) cohort. All Icelandic speaking women, 18–69 years of age, residing in Iceland ($N = 104\,197$ women) were actively recruited in the study. A total of 30 403 women (approximately 30% of eligible women in Iceland) participated in the study. The study population is representative of the total female population in terms of age, education, income, and geographic residence [44]. Participants with missing item(s) on the PSQI were excluded ($n = 722$), which resulted in a study population of 29 681 women (see flowchart in [Supplementary Figure S1](#)).

Procedure

Women were invited to participate in the study through a phone text message or via mail. Participation included answering electronically an extensive internet-based questionnaire on mental and physical morbidities, including sleep disturbances. Data collection took place between March 2018 and July 2019.

Measures

Background information. Demographic information was collected for age (continuous, in years; categorized as: 18–29, 30–39, 40–49, 50–59, 60–69), marital status (married/in a relationship, single/divorced/widowed), number of children (continuous; 0, 1–2, 3–4, ≥ 5), education (primary, secondary, tertiary level A, tertiary level B), personal monthly income (<1200 Euro [EUR], 1201–2400 EUR, 2401–4000 EUR, 4001–5600 EUR, 5601–8000 EUR, >8001 EUR; conversion rates according to the Central Bank of Iceland, February 1, 2018) [45], employment status (active: working/studying/parental leave, inactive: on disability/sick leave/unemployed/retired), work schedule (fixed, flexible, shift work, unemployed), region (Reykjavik capital area, East Iceland, North Iceland, South Iceland, Southern Peninsula, Westfjords, West Iceland, living abroad), and response period (summer [June, July, August, September], fall [October, November], winter [December, January, February, March], spring [April, May]; classified according to the Icelandic Meteorological Office).

Health-related behavior and well-being. Participants answered questions related to their general health, that is, BMI (continuous; categorized as: underweight [<18.5], normal weight [18.5–24.9], overweight [25.0–29.9], obese [≥ 30.0]), smoking (never, previous smoker, non-daily smoker, daily smoker; question based on the Fagerström Test for Nicotine Dependence) [46], alcohol consumption in the past year (i.e. six or more drinks on one occasion [never, less than once a month, monthly, once or more a week; based on an item from the Alcohol Use Disorders Identification Test]) [47], daily screen time during leisure time for the last week (i.e. in front of television, smartphone, and/or computer [<3 h, 3–5 h, 5–7 h, >7 h]). The Patient Health Questionnaire 9-item (PHQ-9) [48] was used to assess depressive symptoms in the past 2 weeks. Items were scored from 0 to 3 for symptom frequency. Standard cut-off scores were used: ≤ 9 indicating no or mild symptoms; 10–14 moderate symptoms; and ≥ 15 moderately

severe/severe symptoms [49]. The General Anxiety Disorder 7-item (GAD-7) [50] questionnaire was used to assess anxiety symptoms in the past 2 weeks. Items were scored on a four-point interval scale ranging from 0 to 3. Symptoms were classified as none/mild (score ≤ 9); moderate (score 10–14); and severe (score ≥ 15) [49].

Sleep quality. The PSQI [3], a 19-item self-report scale, was used to assess sleep problems over a one-month time period. The items were scored on a four-point interval scale ranging from 0 to 3. The items generate seven components' scores, including subjective sleep quality, sleep latency, sleep duration, habitual sleep efficiency, sleep disturbances, use of sleeping medication, and daytime dysfunction. The PSQI global score ranges from 0 to 21, with a lower score indicating better sleep quality, and a score above five indicating clinically significant sleep problems in at least two components or moderate difficulties in more than three components. A total PSQI score of >10 was used as a cut-off for severe sleep problems, in line with previous studies [8]. When analyzing the PSQI components separately, scores ranged from 0 (better) to 3 (worse) and a cut-off score of two was used to indicate symptomology, except for "use of sleep medication," which was classified as present or absent (cut-off score > 1) and for "sleep length," which was classified as sleeping more or less than seven hours (cut-off score > 1). The psychometric properties of the PSQI are adequate, particularly with regard to diagnostic sensitivity (89.6%) and specificity (86.5%) for insomnia [3]. The psychometric properties of the Icelandic version of the scale have demonstrated good internal consistency (i.e. Cronbach's $\alpha = .82$) [51].

Statistics

Descriptive statistics were used to describe the distribution of demographic characteristics, residency, BMI, health-related behavior, well-being, and the seven PSQI components scores of the respondents. We used multiple binary logistic regression model to estimate the prevalence of severe sleep problems. The association of individual variables with severe sleep problems were presented as prevalence ratios (PRs) with 95% confidence intervals (CIs) by utilizing the modified Poisson model with the sandwich variance estimator [52, 53].

We divided the predictors into situational factors and factors related to health-behavior and well-being. A diagram explaining the relationship between the variables is included in the supplement (Supplementary Figure S2). We used multivariable adjustment for the association between situational factors and sleep (i.e. adjustments were made for age [using 10-year intervals], marital status, number of children, education, personal income, work schedule, region, and response period). The variables significantly associated with sleep were used as an adjustment when analyzing the association between severe sleep problems and behavioral factors and severe sleep problems and well-being (i.e. depression and anxiety).

Multiple imputation was used to replace missing values with $m = 20$ rounds of imputations and 20 iterations, using predictive mean matching [54]. The method of Benjamini-Hochberg for false discovery rate was used to correct for multiple comparison. We used a Wald test to test whether there was an effect modification by age groups. Cronbach's α was estimated to determine

internal consistency between the PSQI components. The statistical analyses were done using R (version 4.1.1).

Results

Study cohort

Data from 29 681 women contributed to this analysis. Characteristics of study participants are summarized in Table 1. The mean age of participants was 43.5 years (± 13.7). The majority of the women were married or in a relationship (75.7%), had one or two children (39.3%), a university education (54.0%), 2401–4000 EUR monthly income (31.1%), were active in the labor market (85.2%), had a fixed work schedule (54.1%), and lived in the Reykjavik capital area (66.6%). In addition, 63.2% were overweight or obese, 15.5% were current smokers, and 13.9% binge drink at least once a month. The majority of women (53.0%) spent more than 3 hours every day on leisure-based screen time. Approximately one-third (29.1%) of women had moderate or severe depressive symptoms in the past 2 weeks and 22.6% had moderate or severe anxiety symptoms. Approximately half of the women answered the questionnaire during wintertime (Table 1).

Sleep disturbances

The mean PSQI global score was 7.7 (± 4.0). The majority (65.5%) of the women experienced sleep problems in the past month (PSQI > 5), and nearly a quarter (24.2%) had severe sleep problems (PSQI > 10) (Table 2). Nearly half of the women (47.5%) slept less than seven hours each night in the past month and 44.9% reported sleep latency greater than 30 min at least once a week in the past month. Furthermore, 40.5% of women reported moderate to severe sleep disturbances, for example, waking up in the middle of the night or early morning, coughing or snoring loudly, or feeling too cold or hot. Approximately one-third (32.8%) of women spent $\leq 74\%$ of time asleep while in bed and 32.4% rated their overall sleep quality during the past month as fair or very poor. One-fifth (20.3%) of women had moderate to severe daytime dysfunction and one-third (32.3%) reported using over the counter or prescribed sleep medication in the past month (Table 2; Supplementary Tables S1 and S2).

Social determinants of severe sleep problems

We found a U-shaped association between age and severe sleep problems (PSQI > 10). When assessing the association between individual PSQI components and age we also found a U-shaped relationship between age and both sleep latency > 30 min and moderate/severe difficulty sleeping. In contrast, we found an inverted U-shaped association between age and sleep duration < 7 h. Further, the association between age and both sleep medication use and moderate/severe sleep disturbances was positive, while a negative association was found between age and moderate/severe daytime dysfunction and fairly/very bad subjective sleep quality (Figure 1).

We found being single, divorced, or widowed to be associated with severe sleep problems, compared to women married or in a relationship (adjusted PR [aPR], 1.36; 95% CI = 1.31 to 1.42). Furthermore, compared to having no children, women with one

Table 1. Characteristics of study participants*

	Overall (n = 29 681)	PSQI ≤ 10 (n = 22 484)	PSQI >10 (n = 7197)
Age			
Mean (SD)	43.5 (13.7)	43.4 (13.6)	44.1 (13.9)
Age range			
18–29	5887 (19.8%)	4486 (20.0%)	1401 (19.5%)
30–39	6131 (20.7%)	4744 (21.1%)	1387 (19.3%)
40–49	6554 (22.1%)	5035 (22.4%)	1519 (21.1%)
50–59	6671 (22.5%)	4948 (22.0%)	1723 (23.9%)
60–69	4438 (15.0%)	3271 (14.4%)	1167 (16.2%)
Marital status			
Married or in a relationship	22 359 (75.7%)	17 485 (78.1%)	4874 (68.3%)
Single, divorced, or widowed	7159 (24.3%)	4892 (21.9%)	2267 (31.7%)
Unknown	163	107	56
No. of children			
Mean (SD)	2.1 (1.4)	2.1 (1.4)	2.2 (1.5)
0	5230 (19.5%)	3999 (19.7%)	1231 (19.1%)
1–2	10 537 (39.3%)	8077 (39.7%)	2460 (38.1%)
3–4	9940 (37.1%)	7538 (37.1%)	2402 (37.2%)
≥5	1083 (4.0%)	725 (3.6%)	358 (5.5%)
Unknown	2891	2145	746
Education			
Tertiary level B	6578 (22.3%)	5572 (24.9%)	1006 (14.1%)
Tertiary level A	9378 (31.7%)	7466 (33.3%)	1912 (26.8%)
Secondary	9231 (31.2%)	6689 (29.8%)	2542 (35.6%)
Primary	4369 (14.8%)	2684 (12.0%)	1685 (23.6%)
Unknown	125	73	52
Personal income			
>8001 EUR	1032 (3.6%)	887 (4.1%)	145 (2.1%)
5601–8000 EUR	2901 (10.2%)	2461 (11.4%)	440 (6.4%)
4001–5600 EUR	6780 (23.8%)	5650 (26.2%)	1130 (16.4%)
2401–4000 EUR	8873 (31.1%)	6802 (31.5%)	2071 (30.0%)
1201–2400 EUR	6854 (24.0%)	4301 (19.9%)	2553 (36.9%)
<1200 EUR	2068 (7.3%)	1497 (6.9%)	571 (8.3%)
Unknown	1173	886	287
Employment status†			
Active	25 105 (85.2%)	20 140 (90.1%)	4965 (69.8%)
Inactive	4369 (14.8%)	2219 (9.9%)	2150 (30.2%)
Unknown	207	125	82
Work schedule			
Fixed	15 958 (54.1%)	13 255 (59.3%)	2703 (38.0%)
Flexible	3154 (10.7%)	2471 (11.1%)	683 (9.6%)
Shift work	2809 (9.5%)	2042 (9.1%)	767 (10.8%)
Unemployed	7553 (25.6%)	4591 (20.5%)	2962 (41.6%)
Unknown	207	125	82
Residence			
Reykjavik capital area	19 735 (66.6%)	15 284 (68.1%)	4451 (62.0%)
East Iceland	757 (2.6%)	570 (2.5%)	187 (2.6%)
North Iceland	3087 (10.4%)	2220 (9.9%)	867 (12.1%)
South Iceland	2212 (7.5%)	1631 (7.3%)	581 (8.1%)
Southern Peninsula	1493 (5.0%)	1019 (4.5%)	474 (6.6%)
Westfjords	457 (1.5%)	324 (1.4%)	133 (1.9%)
West Iceland	1456 (4.9%)	1068 (4.8%)	388 (5.4%)
Living abroad	436 (1.5%)	335 (1.5%)	101 (1.4%)
Unknown	48	33	15
Response period			
Summer (June–Sep)	6026 (20.3%)	4753 (21.1%)	1273 (17.7%)
Fall (Oct–Nov)	4207 (14.2%)	3176 (14.1%)	1031 (14.3%)
Winter (Dec–Mar)	15 419 (51.9%)	11 479 (51.1%)	3940 (54.7%)
Spring (Apr–May)	4029 (13.6%)	3076 (13.7%)	953 (13.2%)
Body max index (BMI)			
Mean (SD)	27.9 (6.0)	27.5 (5.8)	29.1 (6.6)
Normal weight (18.5–24.9)	9532 (35.8%)	7698 (38.1%)	1834 (28.6%)
Underweight (<18.5)	275 (1.0%)	204 (1.0%)	71 (1.1%)

Table 1. Continued

	Overall (n = 29 681)	PSQI ≤ 10 (n = 22 484)	PSQI >10 (n = 7197)
Overweight (25.0–29.9)	8635 (32.4%)	6644 (32.9%)	1991 (31.0%)
Obesity (>30)	8194 (30.8%)	5669 (28.0%)	2525 (39.3%)
Unknown	3045	2269	776
Smoking			
Never	12 709 (47.7%)	10 411 (51.5%)	2298 (35.8%)
Previous smoker	9795 (36.8%)	7150 (35.4%)	2645 (41.2%)
Non-daily smoker	1579 (5.9%)	1134 (5.6%)	445 (6.9%)
Daily smoker	2547 (9.6%)	1516 (7.5%)	1031 (16.1%)
Unknown	3051	2273	778
Binge drinking in past year [†]			
Never	12 620 (47.4%)	9643 (47.7%)	2977 (46.5%)
Less than once a month	10 301 (38.7%)	7952 (39.3%)	2349 (36.7%)
Monthly	2704 (10.2%)	1993 (9.9%)	711 (11.1%)
Once or more a week	996 (3.7%)	636 (3.1%)	360 (5.6%)
Unknown	3060	2260	800
Leisure based screen time per day			
<3 h	13 890 (47.0%)	11 105 (49.6%)	2785 (39.1%)
3–5 h	8717 (29.5%)	6451 (28.8%)	2266 (31.8%)
5–7 h	3503 (11.9%)	2475 (11.1%)	1028 (14.4%)
>7 h	3418 (11.6%)	2367 (10.6%)	1051 (14.7%)
Unknown	153	86	67
Depressive symptoms [‡]			
None/mild	18 164 (70.9%)	15 985 (81.1%)	2179 (36.8%)
Moderate	3952 (15.4%)	2429 (12.3%)	1523 (25.7%)
Moderately severe/severe	3503 (13.7%)	1287 (6.5%)	2216 (37.4%)
Unknown	4062	2783	1279
Anxiety symptoms [§]			
None/mild	20 426 (77.4%)	17 049 (84.7%)	3377 (53.8%)
Moderate	3547 (13.4%)	2056 (10.2%)	1491 (23.7%)
Severe	2429 (9.2%)	1016 (5.0%)	1413 (22.5%)
Unknown	3279	2363	916

*Expressed as mean (SD) for continuous variables and proportions for categorical variables. n: sample size.

[†]Active: working, studying or on parental leave; inactive: on disability, sick leave, unemployed, or retired.

[‡]Six or more drinks on one occasion (one drink is defined as simple measure of spirits, one glass of wine or one small beer).

[§]PHQ-9; symptoms past 2 weeks.

[¶]GAD-7; symptoms past 2 weeks.

Table 2. Overall score and components on the PSQI

	Overall (n = 29 681)*
PSQI overall score	
Sleep problems (PSQI > 5)	19 437 (65.5%)
Severe sleep problems (PSQI > 10)	7197 (24.2%)
PSQI components	
1. Fairly or very poor overall sleep quality	9608 (32.4%)
2. Sleep latency > 30 min	13 338 (44.9%)
3. Sleep duration <7 h	14 110 (47.5%)
4. Habitual sleep efficiency ≤ 74%	9734 (32.8%)
5. Moderate to severe sleep disturbances	12 015 (40.5%)
6. Use of sleep medication [†]	9588 (32.3%)
7. Moderate to severe daytime dysfunction	6020 (20.3%)

*Expressed as proportions.

[†]Prescription or over the counter.

or two (aPR, 1.11; 95% CI, 1.03 to 1.20), three or four (aPR, 1.11; 95% CI, 1.02–1.20), or five or more children (aPR, 1.32; 95% CI, 1.18 to 1.47) had higher prevalence of severe sleep problems. Less education was associated with higher prevalence of severe sleep problems as well as low personal income. Being inactive

in the labor market had a strong association with severe sleep problems (aPR, 2.05; 95% CI, 1.96 to 2.14). Having a flexible work schedule (aPR, 1.27; 95% CI, 1.18 to 1.36), working shifts (aPR, 1.47; 95% CI, 1.37 to 1.58), or being unemployed (aPR, 1.98; 95% CI, 1.88 to 2.08) was associated with higher prevalence of severe sleep problems compared to having a fixed work schedule (Figure 2).

Geographic location, seasons, and severe sleep problems

There were some geographic differences in severe sleep problems observed. Compared to living in the Reykjavik capital area, living in Southern Peninsula (aPR, 1.14; 95% CI, 1.06 to 1.23) and North Iceland (aPR, 1.06; 95% CI, 1.00 to 1.13) was associated with higher prevalence of severe sleep problems (Figure 2). Compared to women who participated in the study during the summer, women who responded during the winter (aPR, 1.21; 95% CI, 1.15 to 1.28), fall (aPR, 1.12; 95% CI, 1.04 to 1.20), and spring (aPR 1.09; 95% CI, 1.02 to 1.17) had a higher prevalence of severe sleep problems (Figures 2 and 3). When analyzing the association of PSQI components and response period, we found the prevalence to be highest during winter on all components, except for sleep duration and sleep latency (Supplementary Table S3).




Figure 1. Association between age, severe sleep problems (PSQI > 10), and the PSQI components. The dots represent the proportion of women in each age category experiencing severe sleep problems and lines represent 95% confidence intervals.

Health-related behavior, well-being, and severe sleep problems

Suboptimal health-related behavior (i.e. smoking and binge drinking) and BMI were associated with higher prevalence of severe sleep problems (Figure 4). Particularly, being overweight (aPR, 1.15; 95% CI, 1.09 to 1.21) or obese (aPR, 1.38; 95% CI, 1.31 to 1.45) was associated with higher prevalence of severe sleep problems compared to being normal weight. Daily smokers (aPR,

1.63; 95% CI, 1.53 to 1.73), non-daily smokers (aPR, 1.43; 95% CI, 1.32 to 1.56), and previous smokers (aPR, 1.34; 95% CI, 1.27 to 1.41) had higher prevalence of severe sleep problems compared to women who have never smoked. Moreover, we found that severe sleep problems increased as a function of increased binge drinking. The association between daily leisure-based screen time and severe sleep problems was statistically significant, that is, increased screen time was associated with higher prevalence of severe sleep problems (Figure 4).




Figure 2. Association of demographic characteristics and severe sleep problems during the past month (PSQI > 10)*. *Expressed as proportions and as prevalence ratio (PR) with 95% confidence intervals (CIs). PRs are adjusted for age, marital status, number of children, education, personal income, work schedule, region, and response period.

Compared to women with no or mild depressive symptoms, women with moderate (aRR, 2.91; 95% CI, 2.76 to 3.07) and moderately severe/severe (aRR, 4.46; 95% CI, 4.24 to 4.69) depressive symptoms had a higher prevalence of severe sleep problems.

Similarly, women with moderate (aPR, 2.30; 95% CI, 2.19 to 2.41) and severe (aPR, 3.03; 95% CI, 2.89 to 3.17) anxiety symptoms had a higher prevalence of severe sleep problems, compared to women with none or mild anxiety symptoms (Figure 4).

Additional results

Education, personal income, employment status, work schedule, BMI, smoking, and depressive- and anxiety symptoms were effect modifiers when looking at the association between severe sleep problems and age groups (i.e., young age [18–39],




Figure 3. Association between response period and severe sleep problems (PSQI > 10). The dots represent the proportion of women in each month experiencing severe sleep problems in the preceding month and lines represent 95% confidence intervals.

middle-aged [40–59], and old [60+]). We found that low education, being inactive in the labor market, smoking, and well-being (i.e. symptoms of depression and anxiety) were associated with higher prevalence of severe sleep problems among women of young age compared to those who were middle aged or old. Additionally, low income and working shifts were associated with higher prevalence of severe sleep problems of middle aged and old women compared to those who were young (Supplementary Table S4).

The association between the independent variables and severe sleep problems during the past month (PSQI > 10) were similar using pooled data after multiple imputation (Supplementary Table S5) and complete cases (Supplementary Table S6). In the current study, the internal consistency of the seven PSQI components was acceptable (Cronbach’s $\alpha = .78$).

Discussion

This population-based study among women in Subarctic regions indicates that one in five experience severe sleep problems, particularly during wintertime. We further found U-shaped relationship between age and severe sleep problems, with severe sleep problems being more common among women in their 20s

	PR (95% CI)	
Body max index		
Normal weight	Ref.	■
Underweight	1.12 (0.92-1.36)	■
Overweight	1.15 (1.09-1.21)	■
Obesity	1.38 (1.31-1.45)	■
Smoking		
Never	Ref.	■
Previous	1.34 (1.27-1.41)	■
Non-daily	1.43 (1.32-1.56)	■
Daily	1.63 (1.53-1.73)	■
Binge drinking in past year		
Never	Ref.	■
Less than once a month	1.10 (1.05-1.15)	■
Monthly	1.25 (1.16-1.34)	■
Once or more a week	1.53 (1.40-1.67)	■
Leisure based screen time per day		
< 3 hours	Ref.	■
3 – 5 hours	1.18 (1.12-1.24)	■
5 – 7 hours	1.26 (1.18-1.33)	■
> 7 hours	1.42 (1.34-1.50)	■
Depressive symptoms		
None/mild	Ref.	■
Moderate	2.91 (2.76-3.07)	■
Moderately severe/severe	4.46 (4.24-4.69)	■
Anxiety symptoms		
None/mild	Ref.	■
Moderate	2.30 (2.19-2.41)	■
Severe	3.03 (2.89-3.17)	■

Figure 4. Associations of BMI, health-related behavior, and well-being with severe sleep problems during the past month (PSQI > 10)*. *Expressed as proportions and as prevalence ratio (PR) with 95% confidence intervals (CIs). PRs are adjusted for age, marital status, number of children, education, personal income, work schedule, region, and response period.

and late 50s compared to middle-aged women. Being single, divorced, or widowed, having children, low SES, unemployment, working flexible hours, and shift work were associated with higher prevalence of severe sleep problems. Health-related factors were also associated with severe sleep problems including being overweight or obese, smoking, binge drinking, and excessive leisure-based screen time.

In our study, approximately half of women are not getting the recommended amount of sleep (i.e. at least 7 h) and many experiences prolonged sleep latency. These results are in line with a previous study that found Icelandic adolescents to have delayed bedtimes and shorter sleep duration compared to their European peers [55]. Our results indicate that sleeping problems are more prevalent among Icelandic women than women in other countries with 65% of women experience sleep problems (PSQI > 5), compared to 27%–53% of women in the United States [7], Germany [8], Korea [9], Spain [10], China [11], and Hong Kong [12]. This is possibly due to greater seasonal changes in light exposure and increased exposure to artificial light at night during the short photoperiod of the Arctic winters [20–22, 56]. Indeed, we found that women who reported their sleep quality during the fall and winter had 12%–21% higher prevalence of severe sleep problems compared to women who participated in the summer and also the highest prevalence of severe sleep problems on all PSQI components, except sleep duration. This is consistent with results of a previous study finding that insomnia and fatigue were more common in January than in August among residents of Northern Norway (69°N), while only small seasonal differences in sleep were found among individuals living in Ghana (5°N) [21]. Previous studies have also found seasonal variation in light exposure to be associated with disturbed circadian system [20], rise- and bedtime, and sleep efficiency [21, 22]. In addition, insomnia appears to be more common among women in Subarctic regions [16–18] than women residing elsewhere [19].

We found severe sleep problems to be more prevalent among young and late-midlife women compared to middle-aged women. Previous studies exploring the relationship between age and sleep problems have found conflicting results. While some studies found steady worsening of sleep problems with increasing age, others found sleep problems to increase in late midlife [10, 57]. For instance, it has been indicated that menopause negatively impacts sleep, independent of other factors such as age [58]. Other study has found younger women to report high prevalence of self-reported sleep disturbances [59]. A possible explanation of the discrepancy in prior studies is that the association between sleep and age differs for different aspects of sleep. Indeed, we found that sleep medication uses and sleep disturbances were more common among late-midlife women while prolonged sleep latency, poor subjective sleep quality, and daytime dysfunction were more common among young women.

We found that 36% of women aged 18–29 years reported spending five hours or more in front of screens daily, compared to 20% of women of other ages ($p < .001$). Many factors can negatively impact young adults' sleep, especially the increased social media usage in recent years [60]. Research has shown that smartphones can disrupt sleep through artificial short-waved blue light exposure, which may affect a malfunction of the circadian timing system and melatonin levels [61]. In addition, unhealthy lifestyle, such as inadequate physical activity, alcohol-, and nicotine use can negatively affect sleep [62].

In line with previous research [24, 25], Subarctic women living alone had a higher prevalence of severe sleep problems compared to those in a relationship. This is possibly due to the positive influence of social support and relationships on sleep [63]. Moreover, we found a higher prevalence of severe sleep problems among women who had children. Previous research indicates that parents report more sleep disturbances than childless adults [26]. Interestingly, studies have also found a positive association between children's sleep disruptions and poor sleep quality among parents, regardless of children's age [27].

Overall, we found that socioeconomic hardship and working flexible hours or working shifts were associated with higher prevalence of severe sleep problems among women living in Subarctic regions. These results are consistent with previous studies which have found low SES and unemployment to be associated with higher prevalence of sleep problems [7, 8, 28, 29]. This is possibly due to stress resulting from financial strain, which is associated with both low SES, and sleep problems among women [64]. Indeed, we found that women who were currently inactive in the labor market were more likely to report low income compared to women active in the labor market (62% vs. 25%; $p < .001$). Previous studies have found shift work to be associated with sleep problems [30, 31], especially among women and older adults [65]. Shift work can have a disruptive effect on normal circadian rhythms and result in physiological stress and chronic impairment of cognition [31]. These results indicate the need to target women with low SES and those working shifts when promoting interventions for improved sleep.

Compared to the Reykjavik capital area (64°08.5'N, 21°55.6'V), living in the Southern Peninsula (e.g. Keflavík: 64°00.2'N, 22°33.9'V) was associated with severe sleep problems. Nearly one-third of residents of the Southern Peninsula are experiencing severe sleep problems. This is the region with the greatest socioeconomic- and public health challenges in the country, such as the lowest prevalence of university-educated inhabitants and the highest prevalence of smoking, obesity (BMI >30), and poor physical and mental health [66]; all factors which have been associated with sleep problems. Therefore, further studies assessing the effect of living in Subarctic regions on sleep are needed.

We further found that high BMI and poor health behaviors, such as cigarette smoking, excessive alcohol consumption, and prolonged leisure-based screen time, were associated with higher prevalence of severe sleep problems among women living in Subarctic regions. Previous studies have found obesity to be associated with a higher prevalence of severe sleep problems, compared to being normal weight [8, 32, 33]. This relationship is bidirectional as sleep deprivation can inhibit the production of the hormone leptin which regulates food intake [67] and obesity is a risk factor for the development of obstructive sleep apnea [68]. The relationship between smoking and sleep problems is also well studied on a population level [8, 9, 34, 35]. Further, excessive alcohol consumption has been associated with sleep problems [36, 37], such as sleep continuity and prolonged sleep latency, during the first half of the night, and increased wakefulness, rapid eye movement rebound, and early morning awakenings during the second half of the night [69–71]. Further, our finding that prolonged screen time was associated with severe sleep problems is in line with previous studies suggesting that increased screen time is associated with longer sleep latency, reduced sleep duration, and decreased sleep quality [38–40].

Besides the potential impact of BMI, smoking, binge-drinking, and leisure-based screen time on sleep quality, prior research has also suggested that the onset of disturbed sleep, in turn, can lead to adverse changes in health-related behavior, such as increased risk of alcohol use, smoking, physical inactivity, and overweight or obesity [72].

Consistent with previous research [41], we found a strong association between severe sleep problems and anxiety and depressive symptoms. While insomnia and hypersomnia are symptoms of depression, insomnia has also been found to increase the risk of depressive symptoms, such as suicidal ideation [73, 74]. Therefore, treating sleep problems can be an important step in reducing the risk of subsequent mental health difficulties.

This study has several strengths worth mentioning. First, the SAGA cohort is a large study with over 30 000 participants who are representative of Icelandic women with regard to age, education, income, and residency [44]. It is also the first nationwide, population-based study to investigate a wide range of sleep problems and associated factors among women residing in Subarctic region. Secondly, the measurement of sleep problems used in the study has been validated in different countries, languages, and samples and is a reliable measurement tool.

Several limitations need to be recognized. First, considering that only women between the ages of 18 and 69 were selected for the study, it is not possible to generalize our results to males or to women in other age groups. Second, this study was cross-sectional, making it impossible to infer the causality of the studied associations. Third, there might be significant differences between objective and self-reported measures of sleep, and this study only included the latter.

In conclusion, our results indicate that the prevalence of sleep problems in the general population of Icelandic women is higher than in other international population-based samples of women. We found higher prevalence of severe sleep problems among women responding during the winter months, when sunlight is limited. In addition, we also confirmed that socioeconomic challenges, shift work, suboptimal health behaviors, and mental health problems are also associated with elevated risks of sleep problems among women in Subarctic region of Iceland with high-social welfare. These results are valuable for identifying women at greatest risk of sleep disruptions and would therefore benefit from targeted prevention, in the Subarctic.

Supplementary Material

Supplementary material is available at SLEEP online.

Funding

The SAGA cohort study received funding from the European Research Council (ERC) to UAV (grant agreement no. 726413) and grant of excellence from the Icelandic Research Fund to UAV (grant no. 163362-051). This work was also supported by grant no. 185287-051 to EBT and 163346-053 (to AH) from the Icelandic Research Fund.

Disclosure Statement

None declared.

Acknowledgments

Approval for the study was granted by the Icelandic Bioethics Committee (nr. VSNb2017110046/03.01). The Icelandic Data Protection Authority was notified of the study.

Data Availability

The data underlying this article are not currently publicly available due to data protection laws but may be available from the corresponding author on reasonable request.

References

1. Institute of Medicine Committee on Sleep Medicine and Research. The national academies collection. In: Colten HR, Altevogt BM, eds. *Sleep Disorders and Sleep Deprivation: An Unmet Public Health Problem*. Washington, DC: National Academies Press (US); 2006.
2. Hirshkowitz M, et al. National Sleep Foundation's sleep time duration recommendations: methodology and results summary. *Sleep Health*. 2015;1(1):40–43. doi:10.1016/j.sleh.2014.12.010.
3. Buysse DJ, et al. The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. *Psychiatry Res*. 1989;28(2):193–213. doi:10.1016/0165-1781(89)90047-4.
4. Kwok CS, et al. Self-reported sleep duration and quality and cardiovascular disease and mortality: a dose-response meta-analysis. *J Am Heart Assoc*. 2018;7(15):e008552. doi:10.1161/JAHA.118.008552.
5. Liu Y, et al. Sleep duration and chronic diseases among U.S. adults age 45 years and older: evidence from the 2010 Behavioral Risk Factor Surveillance System. *Sleep*. 2013;36(10):1421–1427. doi:10.5665/sleep.3028.
6. Itani O, et al. Short sleep duration and health outcomes: a systematic review, meta-analysis, and meta-regression. *Sleep Med*. 2017;32:246–256. doi:10.1016/j.sleep.2016.08.006.
7. Beaudreau SA, et al. Validation of the Pittsburgh Sleep Quality Index and the Epworth Sleepiness Scale in older black and white women. *Sleep Med*. 2012;13(1):36–42. doi:10.1016/j.sleep.2011.04.005.
8. Hinz A, et al. Sleep quality in the general population: psychometric properties of the Pittsburgh Sleep Quality Index, derived from a German community sample of 9284 people. *Sleep Med*. 2017;30:57–63. doi:10.1016/j.sleep.2016.03.008.
9. Lee SY, et al. Factors associated with poor sleep quality in the Korean general population: providing information from the Korean version of the Pittsburgh Sleep Quality Index. *J Affect Disord*. 2020;271:49–58. doi:10.1016/j.jad.2020.03.069.
10. Madrid-Valero JJ, et al. Age and gender effects on the prevalence of poor sleep quality in the adult population. *Gac Sanit*. 2017;31(1):18–22. doi:10.1016/j.gaceta.2016.05.013.
11. Tang J, et al. Gender and regional differences in sleep quality and insomnia: a general population-based study in Hunan province of China. *Sci Rep*. 2017;7:43690. doi:10.1038/srep43690.
12. Wong WS, et al. Prevalence of insomnia among Chinese adults in Hong Kong: a population-based study. *J Sleep Res*. 2011;20(1):117–126. doi:10.1111/j.1365-2869.2010.00822.x.
13. Blume C, et al. "How does Austria sleep?" self-reported sleep habits and complaints in an online survey. *Sleep Breath*. 2020;24(2):735–741. doi:10.1007/s11325-019-01982-5.

14. Schlarb AA, et al. Sleep disturbances and mental strain in university students: results from an online survey in Luxembourg and Germany. *Int J Ment Health Syst.* 2017;11(1):24. doi:10.1186/s13033-017-0131-9.
15. Roth T. Insomnia: definition, prevalence, etiology, and consequences. *J Clin Sleep Med.* 2007;3(5 Suppl):S7–S10.
16. Morin CM, et al. Prevalence of insomnia and its treatment in Canada. *Can J Psychiatry.* 2011;56(9):540–548. doi:10.1177/070674371105600905.
17. Ohayon MM, et al. Insomnia and global sleep dissatisfaction in Finland. *J Sleep Res.* 2002;11(4):339–346. doi:10.1046/j.1365-2869.2002.00317.x.
18. Uhlig BL, et al. Prevalence and associated factors of DSM-V insomnia in Norway: the Nord-Trøndelag Health Study (HUNT 3). *Sleep Med.* 2014;15(6):708–713. doi:10.1016/j.sleep.2014.01.018.
19. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders.* 5th ed. Washington, DC: American Psychiatric Association; 2013.
20. Arendt J. Biological rhythms during residence in polar regions. *Chronobiol.* 2012;29(4):379–394. doi:10.3109/07420528.2012.668997.
21. Friberg O, et al. Associations between seasonal variations in day length (photoperiod), sleep timing, sleep quality and mood: a comparison between Ghana (5) and Norway (69). *J Sleep Res.* 2012;21(2):176–184. doi:10.1111/j.1365-2869.2011.00982.x.
22. Friberg O, et al. Sleep timing, chronotype, mood, and behavior at an Arctic latitude (69 N). *Sleep Med Rev.* 2014;15(7):798–807. doi:10.1016/j.sleep.2014.03.014.
23. Wright Kenneth P, et al. Entrainment of the human circadian clock to the natural light-dark cycle. *Curr Biol.* 2013;23(16):1554–1558. doi:10.1016/j.cub.2013.06.039.
24. Troxel WM, et al. Marital/cohabitation status and history in relation to sleep in midlife women. *Sleep.* 2010;33(7):973–981. doi:10.1093/sleep/33.7.973.
25. Whinnery J, et al. Short and long sleep duration associated with race/ethnicity, sociodemographics, and socioeconomic position. *Sleep.* 2014;37(3):601–611. doi:10.5665/sleep.3508.
26. Hagen EW, et al. The sleep-time cost of parenting: sleep duration and sleepiness among employed parents in the Wisconsin Sleep Cohort Study. *Am J Epidemiol.* 2013;177(5):394–401. doi:10.1093/aje/kws246.
27. Varma P, Conduit R, Junge M, Lee VV, Jackson ML. A systematic review of sleep associations in parents and children. *J Child Fam Stud.* 2021;30(9):2276–2288. doi:10.1007/s10826-021-02002-5.
28. Etindele Sosso FA, et al. Influence of socioeconomic status on objective sleep measurement: a systematic review and meta-analysis of actigraphy studies. *Sleep Health.* 2021;7(4):417–428. doi:10.1016/j.sleh.2021.05.005.
29. Wu W, et al. Sleep quality and its associated factors among low-income adults in a rural area of China: a population-based study. *Int J Environ Res.* 2018;15(9):1–18. doi:10.3390/ijerph15092055.
30. Brito RS, et al. Prevalence of insomnia in shift workers: a systematic review. *Sleep Sci.* 2021;14(1):47–54. doi:10.5935/1984-0063.20190150.
31. Marquié J-C, et al. Chronic effects of shift work on cognition: findings from the VISAT longitudinal study. *Occup Environ Med.* 2015;72(4):258–264. doi:10.1136/oemed-2013-101993.
32. Krističević T, et al. The associations between sleep duration and sleep quality with body-mass index in a large sample of young adults. *Int J Environ Res Public Health.* 2018;15(4):758. doi:10.3390/ijerph15040758.
33. Vargas PA, et al. Sleep quality and body mass index in college students: the role of sleep disturbances. *J Am Coll Health.* 2014;62(8):534–541. doi:10.1080/07448481.2014.933344.
34. Cohrs S, et al. Impaired sleep quality and sleep duration in smokers—results from the German Multicenter Study on Nicotine Dependence. *Addict Biol.* 2014;19(3):486–496. doi:10.1111/j.1369-1600.2012.00487.x.
35. Liao Y, et al. Sleep quality in cigarette smokers and nonsmokers: findings from the general population in central China. *BMC Public Health.* 2019;19(1):808. doi:10.1186/s12889-019-6929-4.
36. Ehlers CL, et al. A history of binge drinking during adolescence is associated with poorer sleep quality in young adult Mexican Americans and American Indians. *Psychopharmacology.* 2018;235(6):1775–1782. doi:10.1007/s00213-018-4889-5.
37. Hartwell EE, et al. The association of alcohol severity and sleep quality in problem drinkers. *Alcohol Alcohol.* 2015;50(5):536–541. doi:10.1093/alcac/agn046.
38. Christensen MA, et al. Direct measurements of smartphone screen-time: relationships with demographics and sleep. *PLoS One.* 2016;11(11):e0165331. doi:10.1371/journal.pone.0165331.
39. Exelmans L, et al. Bedtime mobile phone use and sleep in adults. *Soc Sci.* 2016;148:93–101. doi:10.1016/j.socscimed.2015.11.037.
40. Šmotek M, et al. Evening and night exposure to screens of media devices and its association with subjectively perceived sleep: should “light hygiene” be given more attention? *Sleep Health.* 2020;6(4):498–505. doi:10.1016/j.sleh.2019.11.007.
41. Freeman D, et al. Sleep disturbance and psychiatric disorders. *Lancet Psychiat.* 2020;7(7):628–637.
42. Directorate of Health. *Mælaborð: Stuttur svefn fullorðinna. Hlutfall fullorðinna sem sefur að jafnði 6 klukkustundir eða minna á nóttu [Dashboard: Short Sleep for Adults. The Proportion of Adults Who Slept 6 Hours or Less per Night].* <https://www.landlaeknir.is>. Accessed March 31, 2022.
43. Nordic Medico-Statistical Committee, Nordic Social Statistical Committee. *ATC-Code N05C, Sales of Hypnotics and Sedatives by ATC-Code, Year and Country.* http://pxweb.fujitsu.dk/pxweb/en/Nowbase/Nowbase__NOMESCO%2004%20medicine/MED23%20N05%20-%20Hypnotics%20and%20sedatives.px/table/tableViewLayout2/?rxid=c7e95ba0-c0e2-40d3-a2f7-1a49496b9748. Accessed March 31, 2022.
44. Daniélsdóttir HB, et al. Adverse childhood experiences and resilience among adult women: a population-based study. *Elife* 2022;11. doi:10.7554/eLife.71770.
45. Central Bank of Iceland. *Official Exchange Rate.* <https://www.cb.is/statistics/official-exchange-rate/>. Accessed March 31, 2022.
46. Heatherton TF, et al. The Fagerström test for nicotine dependence: a revision of the Fagerstrom Tolerance Questionnaire. *Br J Addict.* 1991;86(9):1119–1127. doi:10.1111/j.1360-0443.1991.tb01879.x.
47. Saunders JB, et al. Development of the Alcohol Use Disorders Identification Test (AUDIT): WHO collaborative project on early detection of persons with harmful alcohol consumption-II. *Addiction.* 1993;88(6):791–804. doi:10.1111/j.1360-0443.1993.tb02093.x.

48. Kroenke K, et al. The PHQ-9: validity of a brief depression severity measure. *J Gen Intern Med.* 2001;16(9):606–613. doi:10.1046/j.1525-1497.2001.016009606.x.
49. Pfizer. Instructions for Patient Health Questionnaire (PHQ) and GAD-7 Measures. <https://www.phqscreeners.com/images/sites/g/files/g10016261/f/201412/instructions.pdf>. Accessed March 31, 2022.
50. Spitzer RL, et al. A brief measure for assessing generalized anxiety disorder: the GAD-7. *Arch Intern Med.* 2006;166(10):1092–1097. doi:10.1001/archinte.166.10.1092.
51. Thordardottir EB, et al. The manifestations of sleep disturbances 16 years post-trauma. *Sleep.* 2016;39(8):1551–1554. doi:10.5665/sleep.6018.
52. Zou GA. modified poisson regression approach to prospective studies with binary data. *Am J Epidemiol.* 2004;159(7):702–706. doi:10.1093/aje/kwh090.
53. Chen W, et al. Comparing performance between log-binomial and robust Poisson regression models for estimating risk ratios under model misspecification. *BMC Med Res Methodol.* 2018;18(1):63. doi:10.1186/s12874-018-0519-5.
54. Van Buuren S. *Flexible Imputation of Missing Data*. 2nd ed. New York: CRC press; 2018:444.
55. Thorleifsdottir B, et al. Sleep and sleep habits from childhood to young adulthood over a 10-year period. *J Psychosom Res.* 2002;53(1):529–537. doi:10.1016/s0022-3999(02)00444-0.
56. Cho Y, et al. Effects of artificial light at night on human health: a literature review of observational and experimental studies applied to exposure assessment. *Chronobiol Int.* 2015;32(9):1294–1310. doi:10.3109/07420528.2015.1073158.
57. Ohayon MM, et al. Meta-analysis of quantitative sleep parameters from childhood to old age in healthy individuals: developing normative sleep values across the human lifespan. *Sleep.* 2004;27(7):1255–1273. doi:10.1093/sleep/27.7.1255.
58. Xu Q, et al. Examining the relationship between subjective sleep disturbance and menopause: a systematic review and meta-analysis. *Menopause.* 2014;21(12):1301–1318. doi:10.1097/GME.0000000000000240.
59. Grandner MA, et al. Age and sleep disturbances among American men and women: data from the US behavioral risk factor surveillance system. *Sleep.* 2012;35(3):395–406. doi:10.5665/sleep.1704.
60. Alonzo R, et al. Interplay between social media use, sleep quality, and mental health in youth: a systematic review. *Sleep Med Rev.* 2021;56:101414. doi:10.1016/j.smrv.2020.101414.
61. Cajochen C, et al. Evening exposure to a light-emitting diodes (LED)-backlit computer screen affects circadian physiology and cognitive performance. *J Appl Physiol (1985).* 2011;110(5):1432–1438. doi:10.1152/jappphysiol.00165.2011.
62. Shochat T. Impact of lifestyle and technology developments on sleep. *Nat Sci Sleep.* 2012;4:19–31. doi:10.2147/NSS.S18891.
63. Kent de Grey RG, et al. Social support and sleep: a meta-analysis. *Health Psychol.* 2018;37(8):787–798. doi:10.1037/hea0000628.
64. Hall MH, et al. Race and financial strain are independent correlates of sleep in midlife women: the SWAN sleep study. *Sleep* 2009;32(1):73–82. doi:10.5665/sleep/32.1.73.
65. Saksvik IB, et al. Individual differences in tolerance to shift work—a systematic review. *Sleep Med Rev.* 2011;15(4):221–235. doi:10.1016/j.smrv.2010.07.002.
66. Directorate of Health. *Súðurnes—Lýðheilsuviðar 2017 [Southern Peninsula—Public Health Indicators 2017]*; 2017. https://www.landlaeknir.is/servelet/file/store93/item35037/Lydheilsuvisar_2018_SUDURNES.pdf
67. Spiegel K, et al. Leptin levels are dependent on sleep duration: relationships with sympathovagal balance, carbohydrate regulation, cortisol, and thyrotropin. *J Clin Endocrinol Metab.* 2004;89(11):5762–5771. doi:10.1210/jc.2004-1003.
68. Romero-Corral A, et al. Interactions between obesity and obstructive sleep apnea: implications for treatment. *Chest.* 2010;137(3):711–719. doi:10.1378/chest.09-0360.
69. Ebrahim IO, et al. Alcohol and sleep I: effects on normal sleep. *Alcohol Clin Exp Res.* 2013;37(4):539–549. doi:10.1111/acer.12006.
70. Fan F, et al. Cohort profile: The Wenchuan Earthquake Adolescent Health Cohort Study. *Int J Epidemiol.* 2017;46(1):27–28. doi:10.1093/ije/dyw013.
71. Thakkar MM, et al. Alcohol disrupts sleep homeostasis. *Alcohol.* 2015;49(4):299–310. doi:10.1016/j.alcohol.2014.07.019.
72. Clark AJ, et al. Onset of impaired sleep as a predictor of change in health-related behaviours; analysing observational data as a series of non-randomized pseudo-trials. *Int J Epidemiol.* 2015;44(3):1027–1037. doi:10.1093/ije/dyv063.
73. Bishop TM, et al. Sleep, suicide behaviors, and the protective role of sleep medicine. *Sleep Med.* 2020;66:264–270. doi:10.1016/j.sleep.2019.07.016.
74. Ford DE, et al. Epidemiologic study of sleep disturbances and psychiatric disorders. An opportunity for prevention? *JAMA* 1989;262(11):1479–1484. doi:10.1001/jama.262.11.1479.